

Versions of the circle method

Edna Jones

Tulane University

Discrete Mathematics Seminar
Kennesaw State University
February 20, 2025

Partitions of a positive integer

Definition (Partition of a positive integer)

Let n be a positive integer. A **partition** of n is a way to write n as the sum of positive integers, where the order of the summands **does not** matter.

Example (Partitions of 5)

$$5$$

$$4 + 1$$

$$3 + 2$$

$$3 + 1 + 1$$

$$2 + 2 + 1$$

$$2 + 1 + 1 + 1$$

$$1 + 1 + 1 + 1 + 1$$

Number of partitions of n

For a positive integer n , let $p(n)$ be the number of partitions of n .
By convention, $p(0) = 1$.

Example (Partitions of 5)

5	$2 + 2 + 1$
$4 + 1$	$2 + 1 + 1 + 1$
$3 + 2$	$1 + 1 + 1 + 1 + 1$
$3 + 1 + 1$	

$$\implies p(5) = 7$$

Number of partitions of n

For a positive integer n , let $p(n)$ be the number of partitions of n .
By convention, $p(0) = 1$.

Example (Partitions of 5)

5	$2 + 2 + 1$
$4 + 1$	$2 + 1 + 1 + 1$
$3 + 2$	$1 + 1 + 1 + 1 + 1$
$3 + 1 + 1$	

$$\implies p(5) = 7$$

How does $p(n)$ grow as $n \rightarrow \infty$?

Generating functions

Definition

The **(ordinary) generating function** for the sequence $(a_n)_{n=0}^{\infty}$ is

$$\sum_{n=0}^{\infty} a_n q^n.$$

Example (Generating function for $(1, 1, 1, \dots)$)

The generating function for the sequence $(1, 1, 1, \dots)$ is

$$1 + q + q^2 + q^3 + \dots = \sum_{k=0}^{\infty} q^k = \frac{1}{1 - q}$$

if $|q| < 1$.

Generating functions

Definition

The **(ordinary) generating function** for the sequence $(a_n)_{n=0}^{\infty}$ is

$$\sum_{n=0}^{\infty} a_n q^n.$$

Example (Generating function for $(p(n))_{n=0}^{\infty}$)

$$\begin{aligned}\sum_{n=0}^{\infty} p(n)q^n &= \prod_{k=1}^{\infty} (q^{0 \cdot k} + q^{1 \cdot k} + q^{2 \cdot k} + q^{3 \cdot k} + \dots) \\ &= \prod_{k=1}^{\infty} \frac{1}{1 - q^k}\end{aligned}$$

for $|q| < 1$.

What is the circle method?

- A collection of techniques for using the analytic properties of the generating function of a sequence to obtain an asymptotic formula for the sequence

What is the circle method?

- A collection of techniques for using the analytic properties of the generating function of a sequence to obtain an asymptotic formula for the sequence
- Typically refers to the Hardy–Littlewood circle method

What is the circle method?

- A collection of techniques for using the analytic properties of the generating function of a sequence to obtain an asymptotic formula for the sequence
- Typically refers to the Hardy–Littlewood circle method
- May also refer to other methods that are used to provide an asymptotic formula for the number of ways an integer is represented by an integer-valued function (like a quadratic form) on \mathbb{Z}^s
 - Kloosterman circle method
 - The delta(-symbol) method

Hardy–Littlewood circle method

- Originally developed by Hardy and Ramanujan (1918) to provide asymptotic formula for the partition function $p(n)$
- Proved that

$$p(n) \sim \frac{1}{4n\sqrt{3}} \exp\left(\pi\sqrt{\frac{2n}{3}}\right)$$

Partition function & modularity

$$f(z) = \sum_{n=0}^{\infty} p(n)e(nz),$$

where $e(z) = e^{2\pi iz}$ and $\text{Im}(z) > 0$.

Partition function & modularity

$$f(z) = \sum_{n=0}^{\infty} p(n)e(nz),$$

where $e(z) = e^{2\pi iz}$ and $\text{Im}(z) > 0$.

$$f(z) = \frac{e(z/24)}{\eta(z)},$$

where $\eta(z)$ is the Dedekind eta function

$$\eta(z) = e\left(\frac{z}{24}\right) \prod_{m=1}^{\infty} (1 - e(mz)).$$

Hardy and Ramanujan used the modularity of η to obtain the asymptotic formula.

Hardy–Littlewood circle method & partition function

$$f(z) = \sum_{n=0}^{\infty} p(n)e(nz)$$

$$F(q) = \sum_{n=0}^{\infty} p(n)q^n$$

where $q = e(z)$.

Hardy–Littlewood circle method & partition function

$$f(z) = \sum_{n=0}^{\infty} p(n)e(nz) \qquad F(q) = \sum_{n=0}^{\infty} p(n)q^n$$

where $q = e(z)$.

Using the Cauchy integral formula, we find that

$$p(n) = \frac{1}{2\pi i} \int_{|q|=r} \frac{F(q)}{q^{n+1}} dq,$$

where $0 < r < 1$.

Hardy–Littlewood circle method & partition function

$$f(z) = \sum_{n=0}^{\infty} p(n)e(nz) \qquad F(q) = \sum_{n=0}^{\infty} p(n)q^n$$

where $q = e(z)$.

Using the Cauchy integral formula, we find that

$$p(n) = \frac{1}{2\pi i} \int_{|q|=r} \frac{F(q)}{q^{n+1}} dq,$$

where $0 < r < 1$.

Changing q into $e(x + iy)$, we obtain

$$p(n) = \int_0^1 f(x + iy)e(-n(x + iy)) dx,$$

where $y > 0$ is such that $r = e^{-2\pi y}$.




Figure: Modulus of $\prod_{m=1}^{\infty} (1 - q^m)$ with $|q| < 1$. From Wikipedia.

Main contribution to integral from points near $e(a/q)$ where q is small ($a, q \in \mathbb{Z}$, $q > 0$).

Major arcs and minor arcs

Split $[0, 1]$ into major arcs \mathfrak{M} and minor arcs \mathfrak{m} .

Major arcs and minor arcs

Split $[0, 1]$ into major arcs \mathfrak{M} and minor arcs \mathfrak{m} .

$$\mathfrak{M} = \left\{ x \in [0, 1] : x \text{ is "close to"} \frac{a}{q}, a, q \in \mathbb{Z}, 0 < q \leq Q \right\}.$$

How close depends on the application of the method.

Major arcs and minor arcs


Split $[0, 1]$ into major arcs \mathfrak{M} and minor arcs \mathfrak{m} .

$$\mathfrak{M} = \left\{ x \in [0, 1] : x \text{ is "close to"} \frac{a}{q}, a, q \in \mathbb{Z}, 0 < q \leq Q \right\}.$$

How close depends on the application of the method.

$$\mathfrak{m} = [0, 1] \setminus \mathfrak{M}.$$

Example of major arcs \mathfrak{M} when $Q = 3$ for the Hardy–Littlewood circle method:



$$\begin{aligned}
 \rho(n) &= \int_0^1 f(x + iy)e(-n(x + iy)) \, dx \\
 &= \underbrace{\int_{\mathfrak{M}} f(x + iy)e(-n(x + iy)) \, dx}_{\text{main term}} + \underbrace{\int_{\mathfrak{m}} f(x + iy)e(-n(x + iy)) \, dx}_{\text{error term}}
 \end{aligned}$$

Real quadratic forms

F is a real quadratic form in s variables \iff
For all $\mathbf{m} \in \mathbb{R}^s$,

$$F(\mathbf{m}) = \frac{1}{2} \mathbf{m}^\top A \mathbf{m},$$

where A is a real symmetric $s \times s$ matrix and is the Hessian matrix of F .

Example (Example of a quadratic form in 2 variables)

$$\begin{aligned} F(\mathbf{m}) &= m_1^2 + m_1 m_2 + m_2^2 \\ &= \frac{1}{2} \mathbf{m}^\top \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} \mathbf{m} \end{aligned}$$

Quadratic form definitions

Definition (Integral quadratic form)

A quadratic form F is **integral** if $F(\mathbf{m}) \in \mathbb{Z}$ for all $\mathbf{m} \in \mathbb{Z}^s$.

Definition (Positive definite quadratic form)

A quadratic form F is **positive definite** if $F(\mathbf{m}) > 0$ for all $\mathbf{m} \in \mathbb{R}^s \setminus \{\mathbf{0}\}$.

Examples (Examples of integral positive definite quadratic forms)

- $f_4(\mathbf{m}) = m_1^2 + m_2^2 + m_3^2 + m_4^2$
- $x^2 + xy + y^2$

Definition ((Unweighted) representation number)

$$R_F(n) = \#\{\mathbf{m} \in \mathbb{Z}^s : F(\mathbf{m}) = n\}$$

Want an asymptotic formula for $R_F(n)$ when F is a positive definite quadratic form.

Definition ((Unweighted) representation number)

$$R_F(n) = \#\{\mathbf{m} \in \mathbb{Z}^s : F(\mathbf{m}) = n\}$$

Want an asymptotic formula for $R_F(n)$ when F is a positive definite quadratic form.

Use same overall method for obtaining an asymptotic formula for the partition function.

Note that the theta function

$$\Theta(z) = \sum_{n=0}^{\infty} R_F(n)e(nz)$$

is a modular form.

Singular series $\mathfrak{S}_F(n)$

$$\mathfrak{S}_F(n) = \prod_p \sigma_{F,p}(n),$$

where $\sigma_{F,p}(n)$ is a p -adic density defined by

$$\sigma_{F,p}(n) = \lim_{k \rightarrow \infty} \frac{\#\{\mathbf{m} \in (\mathbb{Z}/p^k\mathbb{Z})^s : F(\mathbf{m}) \equiv n \pmod{p^k}\}}{p^{(s-1)k}}.$$

An asymptotic for representation numbers from Hardy–Littlewood circle method

Theorem (Kloosterman, 1924)

Suppose that n is a positive integer.

Suppose that F is a positive definite integral quadratic form in $s \geq 5$ variables.

Let $A \in M_s(\mathbb{Z})$ be the Hessian matrix of F .

Then the number of integral solutions to $F(\mathbf{m}) = n$ is

$$R_F(n) = \mathfrak{S}_F(n) \frac{(2\pi)^{s/2}}{\Gamma(s/2) \sqrt{|\det(A)|}} n^{\frac{s}{2}-1} + O_{F,\varepsilon} \left(n^{\frac{s}{4}+\varepsilon} + n^{\frac{s}{2}-\frac{5}{4}+\varepsilon} \right)$$

for any $\varepsilon > 0$.

Motivation for the Kloosterman circle method

- Want a better error term in asymptotic formula for $R_F(n)$ when F is a positive definite quadratic form.
- Split $[0, 1]$ differently.

Farey sequence \mathfrak{F}_Q of order Q

Definition

For $Q \geq 1$, the **Farey sequence \mathfrak{F}_Q of order Q** is the increasing sequence of all reduced fractions $\frac{a}{q}$ with $1 \leq q \leq Q$ and $\gcd(a, q) = 1$.

$$Q = 1$$




Farey sequence \mathfrak{F}_Q of order Q

Definition

For $Q \geq 1$, the **Farey sequence \mathfrak{F}_Q of order Q** is the increasing sequence of all reduced fractions $\frac{a}{q}$ with $1 \leq q \leq Q$ and $\gcd(a, q) = 1$.

$$Q = 2$$




Farey sequence \mathfrak{F}_Q of order Q

Definition

For $Q \geq 1$, the **Farey sequence \mathfrak{F}_Q of order Q** is the increasing sequence of all reduced fractions $\frac{a}{q}$ with $1 \leq q \leq Q$ and $\gcd(a, q) = 1$.

$$Q = 3$$




Farey sequence \mathfrak{F}_Q of order Q


Definition

For $Q \geq 1$, the **Farey sequence \mathfrak{F}_Q of order Q** is the increasing sequence of all reduced fractions $\frac{a}{q}$ with $1 \leq q \leq Q$ and $\gcd(a, q) = 1$.

$$Q = 3$$



Example of Farey dissection when $Q = 3$:



An asymptotic for representation numbers from the Kloosterman method

Theorem

Suppose that n is a positive integer.

Suppose that F is a positive definite integral quadratic form in $s \geq 4$ variables.

Let $A \in M_s(\mathbb{Z})$ be the Hessian matrix of F .

Then the number of integral solutions to $F(\mathbf{m}) = n$ is


$$R_F(n) = \mathfrak{S}_F(n) \frac{(2\pi)^{s/2}}{\Gamma(s/2) \sqrt{|\det(A)|}} n^{\frac{s}{2}-1} + O_{F,\varepsilon} \left(n^{\frac{s-1}{4} + \varepsilon} \right)$$

for any $\varepsilon > 0$.


Kloosterman proved this (with a worse error term) in 1926 for diagonal quadratic forms ($F(\mathbf{m}) = a_1 m_1^2 + \cdots + a_s m_s^2$), using what is now called the Kloosterman circle method.

Hardy–Littlewood vs. Kloosterman

Example of major arcs when $Q = 3$ for the Hardy–Littlewood circle method:



Example of Farey dissection when $Q = 3$ for the Kloosterman circle method:



Hardy–Littlewood vs. Kloosterman

Hardy–Littlewood:

$$R_F(n) = \mathfrak{S}_F(n) \frac{(2\pi)^{s/2}}{\Gamma(s/2)\sqrt{\det(A)}} n^{\frac{s}{2}-1} + O_{F,\varepsilon} \left(n^{\frac{s}{4}+\varepsilon} + n^{\frac{s}{2}-\frac{5}{4}+\varepsilon} \right)$$

Kloosterman:

$$R_F(n) = \mathfrak{S}_F(n) \frac{(2\pi)^{s/2}}{\Gamma(s/2)\sqrt{\det(A)}} n^{\frac{s}{2}-1} + O_{F,\varepsilon} \left(n^{\frac{s-1}{4}+\varepsilon} \right)$$

The delta method

- Rewrite $\delta(n)$, the indicator function for zero, using bump functions

The delta method

- Rewrite $\delta(n)$, the indicator function for zero, using bump functions
- More versatile than Kloosterman circle method

The delta method

- Rewrite $\delta(n)$, the indicator function for zero, using bump functions
- More versatile than Kloosterman circle method
- Developed by Duke, Friedlander, and Iwaniec in 1993 to compute bounds for automorphic L -functions

The delta method

- Rewrite $\delta(n)$, the indicator function for zero, using bump functions
- More versatile than Kloosterman circle method
- Developed by Duke, Friedlander, and Iwaniec in 1993 to compute bounds for automorphic L -functions
- Has been used for a variety of applications, including
 - Asymptotic formulas for weighted representation numbers of quadratic forms (e.g., Heath-Brown, Dietmann, J.)
 - Subconvexity bounds for (twists of) automorphic forms (e.g., Munshi)

(Unweighted) representation number

Definition ((Unweighted) representation number)

$$R_F(n) = \#\{\mathbf{m} \in \mathbb{Z}^s : F(\mathbf{m}) = n\}$$

$$R_F(n) = \sum_{\mathbf{m} \in \mathbb{Z}^s} \mathbf{1}_{\{F(\mathbf{m})=n\}},$$

where $\mathbf{1}_{\{F(\mathbf{m})=n\}}$ is the indicator function

$$\mathbf{1}_{\{F(\mathbf{m})=n\}} = \begin{cases} 1 & \text{if } F(\mathbf{m}) = n, \\ 0 & \text{otherwise.} \end{cases}$$

Bump functions & weighted representation numbers

Definition (Bump function)

The space of real-valued, infinitely differentiable, and compactly supported functions on \mathbb{R}^s is denoted by $C_c^\infty(\mathbb{R}^s)$. A function $\psi \in C_c^\infty(\mathbb{R}^s)$ is called a **bump function**.

Bump functions & weighted representation numbers

Definition (Bump function)

The space of real-valued, infinitely differentiable, and compactly supported functions on \mathbb{R}^s is denoted by $C_c^\infty(\mathbb{R}^s)$. A function $\psi \in C_c^\infty(\mathbb{R}^s)$ is called a **bump function**.

Let $\psi \in C_c^\infty(\mathbb{R}^s)$.

For $X > 0$, define

$$\psi_X(\mathbf{m}) = \psi\left(\frac{1}{X}\mathbf{m}\right).$$

Bump functions & weighted representation numbers

Definition (Bump function)

The space of real-valued, infinitely differentiable, and compactly supported functions on \mathbb{R}^s is denoted by $C_c^\infty(\mathbb{R}^s)$. A function $\psi \in C_c^\infty(\mathbb{R}^s)$ is called a **bump function**.

Let $\psi \in C_c^\infty(\mathbb{R}^s)$.

For $X > 0$, define

$$\psi_X(\mathbf{m}) = \psi\left(\frac{1}{X}\mathbf{m}\right).$$

Definition (Weighted representation number)

$$R_{F,\psi,X}(n) = \sum_{\mathbf{m} \in \mathbb{Z}^s} \mathbf{1}_{\{F(\mathbf{m})=n\}} \psi_X(\mathbf{m})$$

Indicator function

$$\delta(n) = \mathbf{1}_{\{n=0\}} = \begin{cases} 1 & \text{if } n = 0, \\ 0 & \text{otherwise.} \end{cases}$$

$$\delta(n) = \mathbf{1}_{\{n=0\}} = \begin{cases} 1 & \text{if } n = 0, \\ 0 & \text{otherwise.} \end{cases}$$

$$\mathbf{1}_{\{F(\mathbf{m})=n\}} = \delta(F(\mathbf{m}) - n)$$

Indicator function

$$\delta(n) = \mathbf{1}_{\{n=0\}} = \begin{cases} 1 & \text{if } n = 0, \\ 0 & \text{otherwise.} \end{cases}$$

$$\mathbf{1}_{\{F(\mathbf{m})=n\}} = \delta(F(\mathbf{m}) - n)$$

$$\Rightarrow \begin{cases} R_F(n) = \sum_{\mathbf{m} \in \mathbb{Z}^s} \delta(F(\mathbf{m}) - n) \\ R_{F,\psi,X}(n) = \sum_{\mathbf{m} \in \mathbb{Z}^s} \delta(F(\mathbf{m}) - n) \psi_X(\mathbf{m}) \end{cases}$$

The delta method & bump functions

For the delta method, we require $w \in C_c^\infty(\mathbb{R})$, $w(0) = 0$, and $\sum_{q=1}^{\infty} w(q) \neq 0$.

The delta method & bump functions

For the delta method, we require $w \in C_c^\infty(\mathbb{R})$, $w(0) = 0$, and $\sum_{q=1}^{\infty} w(q) \neq 0$.

If n is an integer, then

$$\delta(n) = \frac{1}{\sum_{q=1}^{\infty} w(q)} \sum_{q|n} \left(w(q) - w\left(\frac{|n|}{q}\right) \right),$$

where the sum over $q \mid n$ is taken to be the sum over the positive divisors of n .

The delta method & bump functions

Using the fact that

$$\frac{1}{q} \sum_{a \pmod{q}} e\left(\frac{an}{q}\right) = \begin{cases} 1 & \text{if } q \mid n, \\ 0 & \text{otherwise,} \end{cases}$$

The delta method & bump functions

Using the fact that

$$\frac{1}{q} \sum_{a \pmod{q}} e\left(\frac{an}{q}\right) = \begin{cases} 1 & \text{if } q \mid n, \\ 0 & \text{otherwise,} \end{cases}$$

we have

$$\begin{aligned} \delta(n) &= \frac{1}{\sum_{q=1}^{\infty} w(q)} \sum_{q \mid n} \left(w(q) - w\left(\frac{|n|}{q}\right) \right) \\ &= \frac{1}{\sum_{q=1}^{\infty} w(q)} \sum_{q=1}^{\infty} \frac{1}{q} \sum_{a \pmod{q}} e\left(\frac{an}{q}\right) \left(w(q) - w\left(\frac{|n|}{q}\right) \right) \end{aligned}$$

if n is an integer.

The delta method

$$\delta(n) = \frac{1}{\sum_{q=1}^{\infty} w(q)} \sum_{q=1}^{\infty} \frac{1}{q} \sum_{a \pmod{q}} e\left(\frac{an}{q}\right) \left(w(q) - w\left(\frac{|n|}{q}\right) \right)$$

if n is an integer.

Bump functions are easier to handle analytically than the discontinuous delta function, which helps when analyzing

$$R_F(n) = \sum_{\mathbf{m} \in \mathbb{Z}^s} \delta(F(\mathbf{m}) - n) \quad \text{or}$$

$$R_{F,\psi,\chi}(n) = \sum_{\mathbf{m} \in \mathbb{Z}^s} \delta(F(\mathbf{m}) - n) \psi_{\chi}(\mathbf{m}).$$

Specifics depend on the application of the delta method.

An asymptotic for weighted representation numbers

Theorem (Heath-Brown, 1996)

Suppose that n is an integer.

Suppose that F is a nonsingular integral quadratic form in $s \geq 4$ variables.

Suppose that $\psi \in C_c^\infty(\mathbb{R}^s)$ is a bump function.

Then for $\varepsilon > 0$, the weighted representation number $R_{F,\psi,n^{1/2}}(n)$ is

$$R_{F,\psi,n^{1/2}}(n) = \mathfrak{S}_F(n)\sigma_{F,\psi,\infty}(n, n^{1/2})n^{\frac{s}{2}-1} + O_{F,\psi,s,\varepsilon}\left(n^{\frac{s-1}{4}+\varepsilon}\right),$$

where

$$\sigma_{F,\psi,\infty}(n, X) = \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2\varepsilon} \int_{|F(\mathbf{m}) - \frac{n}{X^2}| < \varepsilon} \psi(\mathbf{m}) \, d\mathbf{m}.$$

Proof uses the delta method with a Kloosterman refinement.

Theorem (J., 2024)

Suppose that n is a positive integer.

Suppose that F is a nonsingular integral quadratic form in $s \geq 4$ variables.

Suppose that $\psi \in C_c^\infty(\mathbb{R}^s)$ is a bump function.

For $\varepsilon > 0$ and sufficiently large X , there is an asymptotic formula for $R_{F,\psi,X}(n)$ where the implicit constants only depend on ψ , s , and ε .

(Other constants dependent on the quadratic form are explicitly computed.)

An asymptotic for weighted representation numbers

Theorem (J., 2024)

Suppose that n is a positive integer.

Suppose that F is a nonsingular integral quadratic form in $s \geq 4$ variables.

Suppose that $\psi \in C_c^\infty(\mathbb{R}^s)$ is a bump function.

For $\varepsilon > 0$ and sufficiently large X , there is an asymptotic formula for $R_{F,\psi,X}(n)$ where the implicit constants only depend on ψ , s , and ε .

(Other constants dependent on the quadratic form are explicitly computed.)

- Used the Kloosterman circle method (and not the delta method)

An asymptotic for weighted representation numbers

Theorem (J., 2024)

Suppose that n is a positive integer.

Suppose that F is a nonsingular integral quadratic form in $s \geq 4$ variables.

Suppose that $\psi \in C_c^\infty(\mathbb{R}^s)$ is a bump function.

For $\varepsilon > 0$ and sufficiently large X , there is an asymptotic formula for $R_{F,\psi,X}(n)$ where the implicit constants only depend on ψ , s , and ε .

(Other constants dependent on the quadratic form are explicitly computed.)

- Used the Kloosterman circle method (and not the delta method)
- If $F(\mathbf{m}) = \frac{1}{2}\mathbf{m}^\top A\mathbf{m}$, then explicit constants depend on the eigenvalues of A and the smallest integer L such that $LA^{-1} \in M_s(\mathbb{Z})$.

An asymptotic for weighted representation numbers

Theorem (J., 2024)

Suppose that n is a positive integer.

Suppose that F is a nonsingular integral quadratic form in $s \geq 4$ variables.

Suppose that $\psi \in C_c^\infty(\mathbb{R}^s)$ is a bump function.

For $\varepsilon > 0$ and sufficiently large X , there is an asymptotic formula for $R_{F,\psi,X}(n)$ where the implicit constants only depend on ψ , s , and ε .

(Other constants dependent on the quadratic form are explicitly computed.)

Explicit constants are used in a variety of applications, including in computations.

Thank you for listening!

My main theorem

Theorem 1.1. *Suppose that n is a positive integer. Suppose that F is a nonsingular integral quadratic form in $s \geq 4$ variables. Let $A \in M_s(\mathbb{Z})$ be the Hessian matrix of F . Let σ_1 be largest singular value of A , and let ν be the number of positive eigenvalues of A . Let L be the smallest positive integer such that $LA^{-1} \in M_s(\mathbb{Z})$. Suppose that $\psi \in C_c^\infty(\mathbb{R}^s)$ is a bump function. Then for $X \geq 1/\sigma_1$ and $\varepsilon > 0$, the weighted representation number $R_{F,\psi,X}(n)$ is*

$$\begin{aligned} & R_{F,\psi,X}(n) \\ &= \mathfrak{S}_F(n) \sigma_{F,\psi,\infty}(n, X) X^{s-2} \\ &+ O_{\psi,s,\varepsilon} \left(\frac{L^{s/2} X^{(s-1)/2+\varepsilon} \sigma_1^{(3-s)/2+\varepsilon}}{\Gamma(\nu/2) \left(\prod_{j=1}^{\nu} \lambda_j \right)^{1/2}} \left(\frac{n}{X^2} - \frac{\rho_\psi^2}{2} \mathbf{1}_{\{\nu>1\}} \sum_{j=\nu+1}^s \lambda_j \right)^{\nu/2-1} \right. \\ &\quad \left. \times \tau(n) \prod_{p|2\det(A)} (1 - p^{-1/2})^{-1} \right) \\ (1.6) \quad &+ O_{\psi,s,\varepsilon} \left(X^{(s-1)/2+\varepsilon} \sigma_1^{(s+1)/2+\varepsilon} L^{s/2} \tau(n) \prod_{p|2\det(A)} (1 - p^{-1/2})^{-1} \right), \end{aligned}$$

where $\lambda_1, \lambda_2, \dots, \lambda_\nu$ are the positive eigenvalues of A and $\lambda_{\nu+1}, \lambda_{\nu+2}, \dots, \lambda_s$ are the negative eigenvalues of A .

Corollary to my main theorem

Corollary 1.5. *Suppose that F is a nonsingular integral quadratic form in $s \geq 4$ variables. Let $A \in M_s(\mathbb{Z})$ be the Hessian matrix of F . Let σ_1 be largest singular value of A , and let ν be the number of positive eigenvalues of A . Let L be the smallest positive integer such that $LA^{-1} \in M_s(\mathbb{Z})$. If n is a positive integer and $\varepsilon > 0$, then the weighted representation number $R_{F,\psi,X}(n)$ is*

$$R_{F,\psi,X}(n) = \mathfrak{S}_F(n) \sigma_{F,\psi,\infty}(n, n^{1/2}) n^{s/2-1} \\ + O_{\psi,s,\varepsilon} \left(\left(\sigma_1^{(s+1)/2+\varepsilon} + \frac{\sigma_1^{(3-s)/2+\varepsilon}}{\Gamma(\nu/2) \left(\prod_{j=1}^{\nu} \lambda_j\right)^{1/2}} \left(1 - \frac{\rho_{\psi}^2}{2} \mathbf{1}_{\{\nu>1\}} \sum_{j=\nu+1}^s \lambda_j\right)^{\nu/2-1} \right) \right. \\ \left. \times n^{(s-1)/4+\varepsilon/2} \tau(n) L^{s/2} \prod_{p|2 \det(A)} (1 - p^{-1/2})^{-1} \right),$$

where $\lambda_1, \lambda_2, \dots, \lambda_{\nu}$ are the positive eigenvalues of A and $\lambda_{\nu+1}, \lambda_{\nu+2}, \dots, \lambda_s$ are the negative eigenvalues of A .

Lemma for Kloosterman circle method

Lemma

Let $f: \mathbb{R} \rightarrow \mathbb{C}$ be a periodic function of period 1 and with real Fourier coefficients (so that $\overline{f(x)} = f(-x)$ for all $x \in \mathbb{R}$). Then

$$\int_0^1 f(x) dx = 2 \operatorname{Re} \left(\sum_{1 \leq q \leq Q} \int_0^{\frac{1}{qQ}} \sum_{\substack{Q < d \leq q+Q \\ qdx < 1 \\ \gcd(d,q)=1}} f\left(x - \frac{d^*}{q}\right) dx \right),$$

where d^* is the multiplicative inverse of d modulo q .

Use this for

$$f(x) = \sum_{\mathbf{m} \in \mathbb{Z}^s} e((x + iy)(F(\mathbf{m}) - n)),$$

where $y > 0$.